

COHOMOGENEITY-TWO TORUS ACTIONS ON NON-NEGATIVELY CURVED MANIFOLDS OF LOW DIMENSION

FERNANDO GALAZ-GARCIA AND MARTIN KERIN*

ABSTRACT. Let M^n , $n \in \{4, 5, 6\}$, be a compact, simply connected n -manifold which admits some Riemannian metric with non-negative curvature and an isometry group of maximal possible rank. Then any smooth, effective action on M^n by a torus T^{n-2} is equivariantly diffeomorphic to an isometric action on a normal biquotient. Furthermore, it follows that any effective, isometric circle action on a compact, simply connected, non-negatively curved four-dimensional manifold is equivariantly diffeomorphic to an effective, isometric action on a normal biquotient.

The classification of (compact) Riemannian manifolds (M^n, g) with positive or non-negative (sectional) curvature is a notoriously difficult problem. One of the few successes in this quest occurs when one considers such manifolds equipped with an effective action by a suitably large group G of isometries. The ambiguity of the term “suitably large” allows various classification results to be achieved (see, for example, [BB, GS1, GS2, GWZ, Ve, Wa, Wi1, Wi2] and the surveys [Gr, Wi3]).

There are, in fact, two parts to the classification program. First is the *topological classification*, the goal of which is to determine, up to diffeomorphism, all possible positively or non-negatively curved manifolds on which G can act. The second part is the *equivariant classification*, where the goal is to determine all possible actions of G on a given positively or non-negatively curved manifold up to equivariant diffeomorphism.

This approach to the classification problem is inspired by the work of Hsiang and Kleiner [HK]. They showed that a simply connected, four-dimensional manifold (M^4, g) with positive curvature admitting an effective, isometric circle action must be homeomorphic to either S^4 or to \mathbb{CP}^2 . If (M^4, g) is assumed to have only non-negative curvature then, by Kleiner [Kl] and Searle and Yang [SY], M^4 must be homeomorphic to one of S^4 , \mathbb{CP}^2 , $\mathbb{CP}^2 \# \pm \mathbb{CP}^2$ or $S^2 \times S^2$. In both situations, homeomorphism is improved to diffeomorphism by appealing to work of Fintushel [Fi1, Fi2], Pao [Pao] and Perelman’s proof of the Poincaré conjecture [Pe1, Pe2].

The existence of an effective, isometric circle action is equivalent to the rank of the isometry group $\text{Iso}(M^n, g)$ being positive. The success in dimension four suggests that it may be beneficial to consider the topological classification when “largeness” of our group G of isometries refers to the *symmetry rank* of (M^n, g) , defined as the rank of $\text{Iso}(M^n, g)$ and denoted $\text{symrank}(M^n, g)$. Indeed, Grove and Searle [GS1] showed that if (M^n, g) is positively curved, then $\text{symrank}(M^n, g) \leq \lfloor \frac{n+1}{2} \rfloor$. Moreover, if the symmetry

Date: November 8, 2011.

2000 Mathematics Subject Classification. 53C20.

Key words and phrases. non-negative curvature, circle action, torus action, 4-manifolds, 5-manifolds, biquotients, Lie groups, symmetry rank, cohomogeneity.

* This research was carried out as part of SFB 878: Groups, Geometry & Actions, at the University of Münster.

rank is maximal, namely $\text{symrank}(M^n, g) = \lfloor \frac{n+1}{2} \rfloor$, then M^n is diffeomorphic to a sphere, a real or complex projective space, or a lens space.

In the non-negative curvature setting, when (M^n, g) is simply connected and $n \leq 9$, it is known that $\text{symrank}(M^n, g) \leq \lfloor \frac{2n}{3} \rfloor$. If equality holds, then M^n is diffeomorphic to S^4 , \mathbb{CP}^2 , $\mathbb{CP}^2 \# \pm \mathbb{CP}^2$ or $S^2 \times S^2$ for $n = 4$; to $S^3 \times S^2$, $S^3 \tilde{\times} S^2$ (the non-trivial S^3 -bundle over S^2) or S^5 for $n = 5$; and to $S^3 \times S^3$ for $n = 6$ (cf. [GGS]).

This article is concerned with the equivariant classification in low dimensions. Before the statement of the main result, it is necessary to recall that a *biquotient* is a quotient of a Lie group G by the two-sided, free action of a subgroup $U \subset G \times G$. If G is equipped with a bi-invariant metric, then the action of U is by isometries and the quotient G/U equipped with the induced metric (of non-negative curvature) is called a *normal biquotient*.

Theorem A. *Let M^n , $n \in \{4, 5, 6\}$, be an n -manifold which admits a Riemannian metric with non-negative curvature and maximal symmetry rank. Then every smooth, effective action on M^n by a torus T^{n-2} is equivariantly diffeomorphic to an effective, isometric action on a normal biquotient.*

In light of the topological classification discussed earlier, we make the following remarks. Any smooth, effective T^4 action on $S^3 \times S^3$ is equivariantly diffeomorphic to the standard action of T^4 on the Lie group $S^3 \times S^3$ equipped with a bi-invariant metric (cf. [Oh1]). From [Oh2] it follows that any smooth, effective T^3 action on S^5 is equivariantly diffeomorphic to a linear action (on a normal homogeneous space). Similarly, by the classification result of Orlik and Raymond [OR1], any smooth, effective T^2 action on S^4 or \mathbb{CP}^2 is equivariantly diffeomorphic to a linear action (on a normal homogeneous space).

In order to establish Theorem A, it therefore remains only to consider the actions of maximal rank tori on the manifolds $\mathbb{CP}^2 \# \pm \mathbb{CP}^2$, $S^2 \times S^2$, $S^3 \times S^2$ and $S^3 \tilde{\times} S^2$. Each of these manifolds can be described as a normal biquotient of $S^3 \times S^3$ by the free action of either a two-torus or circle (cf. [Ch], [To], [Pav], [DV]). The standard effective, isometric action of T^4 on the Lie group $S^3 \times S^3$ induces a maximal rank, effective, isometric torus action on each of these biquotients. An examination of all possible induced actions on these biquotients then shows that every equivariant diffeomorphism type appearing in the classifications of Orlik and Raymond [OR1] and Oh [Oh2] is achieved.

Grove and Wilking [GW] have recently shown that an effective, isometric circle action on S^4 or \mathbb{CP}^2 is equivariantly diffeomorphic to a linear action. Furthermore, any effective, isometric circle action on a simply connected 4-manifold with non-negative curvature extends to a smooth, effective T^2 action (cf. [GG], [GW]). This fact, in combination with Theorem A, can be applied to $\mathbb{CP}^2 \# \pm \mathbb{CP}^2$ and $S^2 \times S^2$ to draw the following conclusion.

Corollary B. *An effective, isometric circle action on a non-negatively curved, simply connected 4-manifold is equivariantly diffeomorphic to an effective, isometric action on a normal biquotient.*

It is worthwhile to remark that there exist smooth, effective (non-isometric) circle actions on S^4 which do not extend to a smooth T^2 action (cf. [Pao]).

The paper is divided as follows. In Section 1 definitions and notation necessary for the rest of the paper are gathered. In Section 2 some general facts about smooth, cohomogeneity-two torus actions are recalled, while in Sections 3 and 4 all possible orbit spaces of smooth, cohomogeneity-two torus actions on the manifolds $\mathbb{CP}^2 \# \pm \mathbb{CP}^2$,

$S^2 \times S^2$, $S^3 \times S^2$ and $S^3 \tilde{\times} S^2$ are detailed. In Section 5 descriptions of these manifolds as biquotients are given and in Sections 6 and 7 it is shown that isometric, cohomogeneity-two actions on these biquotients recover all of the orbit spaces described in Sections 3 and 4, thus proving Theorem A. Finally, in Section 8 we make some observations about principal circle bundles over the 4-manifolds $\mathbb{CP}^2 \# \pm \mathbb{CP}^2$ and $S^2 \times S^2$.

Acknowledgments. The first named author thanks B. Wilking and K. Grove for useful conversations. The second named author wishes to thank J. DeVito for several interesting and helpful discussions.

1. BASIC DEFINITIONS AND NOTATION

Let $G \times M \rightarrow M$, $m \mapsto g \star m$, be a smooth action of a compact Lie group G on a smooth manifold M . We will denote the orbit space M / G of the action by M^* . The *cohomogeneity* of the action is the dimension of the orbit space M^* . The orbit $G \star p$ through a point $p \in M$ is diffeomorphic to the quotient G / G_p , where $G_p := \{g \in G \mid g \star p = p\}$ is the isotropy subgroup of G at p . If G_p acts trivially on the normal space to the orbit at p , then G / G_p is called a *principal orbit*. The set P of principal orbits is open and dense in M . In fact, the isotropy groups of principal orbits are all conjugate in G , hence each principal orbit has the same dimension.

If G / G_p has the same dimension as a principal orbit, but G_p acts non-trivially on the normal space at p , then G / G_p is called an *exceptional orbit*. An orbit that is neither principal nor exceptional is called a *singular orbit*. The set of exceptional orbits will be denoted by E and the set of singular orbits by Q . When $G_p = G$, p is called a *fixed point* of the action. The set of fixed points of the action will be denoted by $\text{Fix}(M, G)$.

Given a subset $X \subset M$, we will denote its projection under the orbit map $\pi : M \rightarrow M^*$ by X^* . Given a subset $X^* \subset M^*$, we will let $X = \pi^{-1}(X^*)$ be its pre-image under π . Recall that the *ineffective kernel* of the action is $\text{Ker} := \{g \in G \mid g \star m = m, \forall m \in M\}$. The action is *effective* if the ineffective kernel is trivial. The group $\tilde{G} := G / \text{Ker}$ will always act effectively. We say that the action of G is *free* if $g \star m = m$ for some $m \in M$ implies that $g = e$, the identity in G . In this case the orbit space M / G is a manifold. We may expand our definition of freeness to allow an ineffective kernel, namely $g \star m = m$ for some $m \in M$ implies that $g \star m = m$ for all $m \in M$. In this case, the orbit space M / G is again a manifold, diffeomorphic to M / \tilde{G} , where $\tilde{G} = G / \text{Ker}$ acts freely and effectively on M . If, in addition, there is a metric on M such that the action of G is by isometries, then there is an induced metric on M / G , given by the distance between orbits in M .

Let $a_1, \dots, a_n \in \mathbb{Z}$ be relatively prime integers. Define the circle subgroup of slope $\underline{a} = (a_1, \dots, a_n) \in \mathbb{Z}^n$ in T^n via $G(\underline{a}) = G(a_1, \dots, a_n) := \{z^{a_1}, \dots, z^{a_n} \mid z \in \mathbb{C}, |z|^2 = 1\}$.

By the *determinant of n circle subgroups* $G(\underline{a}_1), \dots, G(\underline{a}_n)$ of T^n we mean the determinant of the $(n \times n)$ -matrix

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix}$$

where $\underline{a}_i = (a_{i1}, \dots, a_{in}) \in \mathbb{Z}^n$.

We recall that two subgroups $G(\underline{a}_1)$ and $G(\underline{a}_2)$ of T^n have trivial intersection if and only if there exist $G(\underline{a}_3), \dots, G(\underline{a}_n) \subset T^n$ such that the determinant of these n circle

subgroups is ± 1 . A collection of n circle subgroups $G(\underline{a}_i)$, $i = 1, \dots, n$, span T^n provided their determinant is non-zero, i.e. the vectors $\underline{a}_1, \dots, \underline{a}_n$ span \mathbf{R}^n . They are generators of T^n , that is, $T^n = G(\underline{a}_1) \times \dots \times G(\underline{a}_n)$, if and only if their determinant is ± 1 .

Consider now a closed subgroup $U \subset G \times G$ acting on a compact Lie group G via

$$(u_1, u_2) \star g = u_1 g u_2^{-1}, \quad g \in G, (u_1, u_2) \in U.$$

This action is free if and only if, for non-trivial $(u_1, u_2) \in U$, u_1 is never conjugate to u_2 in G . The resulting quotient manifold, denoted $G // U$, is called a *biquotient*. As discussed above, we may extend the definition of biquotient to allow actions which act freely up to an ineffective kernel (cf. [Es]). In particular, if G is equipped with a bi-invariant metric $\langle \cdot, \cdot \rangle_0$, then U acts by isometries and $(G, \langle \cdot, \cdot \rangle_0) // U$ is called a *normal biquotient*.

2. COHOMOGENEITY-TWO TORUS ACTIONS

Let M^{n+2} be a smooth, compact, simply connected $(n+2)$ -manifold on which a compact, connected Lie group G acts smoothly, effectively and with cohomogeneity two. It is well-known (see, for example, [Br, Chapter IV]) that in the presence of singular orbits, the orbit space M^* of the action is homeomorphic to a 2-disk D^2 with boundary Q^* , the projection of the singular orbits, while the interior points of M^* correspond to principal orbits. When $G = T^n$, $n \geq 2$, the orbit space structure was analyzed in [OR1, KMP] (see also [Oh2]). In this case the only possible non-trivial isotropy groups are S^1 and T^2 . The boundary circle, Q^* , is a union of arcs, and the interior of each arc corresponds to orbits with isotropy S^1 , while the endpoints of each arc correspond to orbits with isotropy T^2 (see Figure 1). Moreover, there must be at least n orbits with isotropy T^2 .




Figure 1: Orbit space of a T^n action on M^{n+2}

Suppose, on the other hand, that $n \geq 2$ and M^* is a two-dimensional manifold homeomorphic to D^2 . Partition the boundary of M^* into N (ordered) arcs ($N \geq n$) and to each arc assign an n -tuple $\underline{x}_i = (x_{i1}, \dots, x_{in}) \in \mathbf{Z}^n$, $i = 1, \dots, N$, where $\gcd(x_{i1}, \dots, x_{in}) = 1$. Consider each of these n -tuples \underline{x}_i to be the slope of a circle $G(\underline{x}_i)$ in T^n . We say that M^* is *legally weighted* with weights $\{\underline{x}_1, \dots, \underline{x}_N\}$ if, for any two adjacent slope vectors \underline{x}_i and \underline{x}_{i+1} (i cyclic index), there exist $n-2$ other n -tuples in $\underline{v}_1, \dots, \underline{v}_{n-2} \in \mathbf{Z}^n$ such that the $(n \times n)$ -matrix with rows $\underline{x}_i, \underline{x}_{i+1}, \underline{v}_1, \dots, \underline{v}_{n-2}$ has determinant ± 1 . This is equivalent to saying that the circles $G(\underline{x}_i)$ and $G(\underline{x}_{i+1})$ have trivial intersection, that is, there exist $n-2$ other circles in T^n such that together the circles generate (the homology of) T^n . By [Oh2], for any legally weighted, orientable 2-manifold M^* there is an $(n+2)$ -dimensional manifold M with orbit space M^* under an effective T^n action.

In fact, following similar techniques to those used in [OR1], Oh [Oh2] showed that two $(n + 2)$ -manifolds M and M' with smooth, effective T^n actions are equivariantly diffeomorphic if and only if there is a weight-preserving diffeomorphism between their orbit spaces M^* and $(M')^*$.

Therefore, given an $(n + 2)$ -manifold M equipped with a smooth, effective T^n action, we may represent M uniquely in terms of its weighted orbit space:

$$M^{n+2} = \{\underline{x}_1, \dots, \underline{x}_N\}, \quad N \geq n \geq 2,$$

where $\underline{x}_1, \dots, \underline{x}_N \in \mathbf{Z}^n$ and $\gcd(\underline{x}_i) = 1$, for $i = 1, \dots, N$.

Moreover, [Oh2] has used the (legally) weighted orbit space M^* to give information about the fundamental group of the manifold M . Indeed, if M^* is a 2-dimensional disk and there are no exceptional orbits (i.e. no finite isotropy groups), and if $\underline{x}_i \in \mathbf{Z}^n$ denotes the slope of the i^{th} S^1 isotropy group $G(\underline{x}_i) \subset T^n$, $i = 1, \dots, N$, then

$$(2.1) \quad \pi_1(M) \subset \mathbf{Z}^n / \langle \underline{x}_1, \dots, \underline{x}_N \rangle.$$

In particular, if the S^1 isotropy groups span T^n (i.e. some n of the \underline{x}_i give an $(n \times n)$ -matrix with nonzero determinant), then $\pi_1(M)$ is finite, and if the S^1 isotropy groups generate T^n (i.e. some n of the \underline{x}_i give an $(n \times n)$ -matrix with determinant ± 1), then M is simply connected. Note that, if the S^1 isotropy groups do not span T^n , then M is diffeomorphic to a product $M' \times S^1$, where M' is an $(n + 1)$ -manifold with an effective T^{n-1} action (cf. [Oh2]). Hence we will always assume that the S^1 isotropy groups span T^n .

That the S^1 isotropy groups give trivial elements of the fundamental group can be explained as follows. Let $p \in M$ such that $T^n \star p$ is a principal orbit and let $q \in M$ such that $T^n \star q$ is a singular orbit with isotropy group $G(\underline{x}_i)$ of slope \underline{x}_i . Take a geodesic $\gamma_i(t)$ from p to q . Then $G(\underline{x}_i) \star \gamma_i(t)$ is a disk, since $G(\underline{x}_i) \star q = q$, and hence $G(\underline{x}_i) \star p$ is homotopically trivial.

3. T^2 ACTIONS ON 4-MANIFOLDS

As discussed in Section 2, the orbit space M^* of a smooth, effective T^2 action on a smooth, compact, simply connected 4-manifold M is an oriented 2-disk whose interior consists of principal orbits and boundary contains $k \geq 2$ isolated points corresponding to orbits with isotropy T^2 , i.e. fixed points. Arcs in the boundary between these points correspond to orbits with circle isotropy groups $G(m_i, n_i)$, $1 \leq i \leq k$, where $\gcd(m_i, n_i) = 1$.

Represent M in terms of its weighted orbit space, that is

$$M = \{\underline{x}_1 = (m_1, n_1), \dots, \underline{x}_k = (m_k, n_k)\}.$$

In order for an orbit space M^* to be legally weighted, it is necessary to have

$$(3.1) \quad \det \begin{pmatrix} m_k & n_k \\ m_1 & n_1 \end{pmatrix} = \varepsilon_1 = \pm 1 \quad \text{and} \quad \det \begin{pmatrix} m_{i-1} & n_{i-1} \\ m_i & n_i \end{pmatrix} = \varepsilon_i = \pm 1, \quad i = 2, 3, \dots, k.$$

Assume without loss of generality that M^* is oriented from $\underline{x}_i = (m_i, n_i)$ to $\underline{x}_{i+1} = (m_{i+1}, n_{i+1})$ and recall that actions on M are in one-to-one correspondence with weights in the orbit space.

Orlik and Raymond [OR1] have shown that M is equivariantly diffeomorphic to a connected sum of S^4 , $\pm \mathbf{CP}^2$ and $S^2 \times S^2$. However, as we are only interested in 4-manifolds which admit a Riemannian metric of non-negative curvature and maximal

symmetry rank, in particular $\mathbf{CP}^2 \# \pm \mathbf{CP}^2$ and $\mathbf{S}^2 \times \mathbf{S}^2$, we may restrict our attention to the case where there are exactly four isolated fixed points on the boundary. This follows from the fact that $\chi(M) = \chi(\text{Fix}(M, T^2))$.

Without loss of generality we may reparametrize our T^2 action and assume that the weighted orbit space is of the form shown in Figure 2.




Figure 2: Weighted orbit space for a T^2 action on M^4

Since the orbit space is assumed to be legally weighted, it follows from (3.1) that $\varepsilon_2 = 1$, $a = -\varepsilon_3$, $d = -\varepsilon_1$ and hence $\varepsilon_1\varepsilon_3 - bc = \varepsilon_4 = \varepsilon_2\varepsilon_4$. Therefore $ad = -2\varepsilon_4$ (whenever $\varepsilon_1\varepsilon_3 = -\varepsilon_2\varepsilon_4 = -\varepsilon_4$) or 0 (whenever $\varepsilon_1\varepsilon_3 = \varepsilon_2\varepsilon_4 = \varepsilon_4$), from which it follows that $(b, c) = \pm(1, -2\varepsilon_4)$ or $\pm(-2\varepsilon_4, 1)$ or $(0, k)$ or $(k, 0)$ for some $k \in \mathbf{Z}$. From [OR1, Section 5], in the first case M is diffeomorphic to $\mathbf{CP}^2 \# \mathbf{CP}^2$, while in the latter two cases M is diffeomorphic to $\mathbf{CP}^2 \# -\mathbf{CP}^2$ or $\mathbf{S}^2 \times \mathbf{S}^2$, depending on the parity of k . Thus, up to reparametrization and reordering, all possible legally weighted orbit spaces are given in Figure 3 below.




Figure 3: Possible legally weighted orbit spaces for $\mathbf{CP}^2 \# \pm \mathbf{CP}^2$ and $\mathbf{S}^2 \times \mathbf{S}^2$

4. T^3 ACTIONS ON 5-MANIFOLDS

Let M be a smooth, compact, simply connected 5-manifold with a smooth, effective T^3 action. By the discussion in Section 2, the orbit space M^* of the action is an oriented 2-disk whose interior consists of principal orbits and its boundary contains $k \geq 3$ isolated points corresponding to orbits with isotropy T^2 . As before, arcs in the boundary between these points correspond to orbits with circle isotropy groups $G(\ell_i, m_i, n_i)$, $2 \leq i \leq k$, where

$\gcd(\ell_i, m_i, n_i) = 1$, and M may be represented in terms of its weighted orbit space, i.e.

$$M = \{\underline{x}_1 = (\ell_1, m_1, n_1), \dots, \underline{x}_k = (\ell_k, m_k, n_k)\}.$$

Oh [Oh2] has classified such manifolds up to equivariant diffeomorphism. Indeed, if $k = 3$, M (together with its T^3 action) is equivariantly diffeomorphic to S^5 equipped with a linear action, while, for $k \geq 4$, M is equivariantly diffeomorphic to either $\#(k-3)(S^3 \times S^2)$ or $(S^3 \tilde{\times} S^2) \# (k-4)(S^3 \times S^2)$, depending respectively on whether the second Stiefel-Whitney class $w_2(M)$ is trivial or not.

As manifolds which admit a Riemannian metric with non-negative curvature and maximal symmetry rank are the focus of our attention, by [GGS] we may therefore assume that $k = 4$, namely that M is one of $S^3 \times S^2$ or $S^3 \tilde{\times} S^2$.

By a suitable reparametrization of the T^3 action, if necessary, we may assume that the orbit space of these manifolds looks like the one in Figure 4 below.





Figure 4: Weighted orbit space for a T^3 action on M^5

Notice that $\underline{x}_1 = (1, 0, 0)$, $\underline{x}_2 = (0, 1, 0)$, $\underline{x}_3 = (p, q, r)$ and $\underline{x}_4 = (x, y, z)$ generate $\langle (1, 0, 0), (0, 1, 0), (0, 0, \gcd(r, z)) \rangle \subset \mathbf{Z}^3$. Therefore, from (2.1) we deduce that $\pi_1(M) \subset \mathbf{Z}_{\gcd(r, z)}$. In fact, as the following proposition shows, we can do better.

Proposition 4.1. *If T^3 acts smoothly and effectively on a smooth, compact 5-manifold M with (legally) weighted orbit space as in Figure 4, then $\pi_1(M) = \mathbf{Z}_{\gcd(r, z)}$. In particular, if M is simply connected, then $\gcd(r, z) = 1$.*

Proof. We want to use the theorem of van Kampen. Let $b \in M$ lie on a principal orbit and let $\{\underline{e}_1, \underline{e}_2, \underline{e}_3\}$ denote the standard basis in \mathbf{R}^3 . The circle of slope (ℓ_1, ℓ_2, ℓ_3) in T^3 is, as always, denoted by $G(\ell_1, \ell_2, \ell_3)$. Set $G_i := G(\underline{e}_i)$, $i = 1, 2, 3$. In particular, from [Oh2] (cf. (2.1)) we know that $\pi_1(M, b)$ is generated by the homotopy classes (of circle orbits) $[G_1 \star b]$, $[G_2 \star b]$ and $[G_3 \star b]$. Denote the arcs in M^* with weights $(0, 1, 0)$ and (x, y, z) by L_1^* and L_2^* respectively, and their respective pre-images under the T^3 action by L_1 and L_2 .

Divide M into two open regions Ω_1 and Ω_2 , centred around L_1 and L_2 respectively, and with $b \in \Omega_1 \cap \Omega_2$. We may assume that Ω_i is a 2-disk bundle over the corresponding arc L_i , $i = 1, 2$, since the isotropy group G_3 (resp. $G(x, y, z)$) acts freely away from L_1 (resp. L_2) (see [Br, Chapter VI, Theorem 2.2]). Then the orbit space of M under the T^3 action is as in Figure 5.

Figure 5: Van Kampen decomposition of M

The points in the interior of the arc L_1^* correspond to points in M with isotropy G_2 , while the endpoints of L_1^* correspond to points in M with isotropy given by $G_1 \times G_2$ and $G_2 \times G(p, q, r)$ respectively. We can therefore think of L_1^* as the quotient of a $G_1 \times G_3 = T^3 / G_2$ action on the 3-manifold L_1 , where the interior of L_1^* corresponds to points with trivial isotropy and the endpoints to points with isotropy $G(1, 0)$ and $G(p, r)$ respectively. By the work of Orlik and Raymond [OR1], we know that L_1 is a lens space $L(r; p)$.

As Ω_1 is a D^2 -bundle over L_1 , and since $G_1 \star b$ and $G_2 \star b$ are homotopically trivial in Ω_1 , it follows that $\pi_1(\Omega_1, b) \cong \pi_1(L_1) = \langle [G_3 \star b] \rangle / \langle p[G_3 \star b] \rangle = \mathbf{Z}_r$.


Since M^* is legally weighted, there exist $\lambda, \mu \in \mathbf{Z}$ such that $\lambda y + \mu z = 1$. Let $A : \mathbf{R}^3 \rightarrow \mathbf{R}^3$ be the linear map given by

$$A = \begin{pmatrix} 1 & -\lambda x & -\mu x \\ 0 & z & -y \\ 0 & \lambda & \mu \end{pmatrix}.$$

Since $\det A = 1$, it follows that we can reparametrize our T^3 action by A , that is, $A\mathbf{e}_i$, $i = 1, 2, 3$, generate T^3 . Notice in particular that

$$A\mathbf{e}_1 = \mathbf{e}_1, \quad A \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \mathbf{e}_3 \quad \text{and} \quad A \begin{pmatrix} p \\ q \\ r \end{pmatrix} = \begin{pmatrix} p - (\lambda q + \mu r)x \\ qz - ry \\ \lambda q + \mu r \end{pmatrix}.$$

In terms of the new parameters, denote circles of slope \mathbf{e}_i and $(p - (\lambda q + \mu r)x, qz - ry, \lambda q + \mu r)$ by \tilde{G}_i and \tilde{G}' respectively. Then we may relabel Ω_2^* as in Figure 6 below.

Figure 6: Ω_2^* after reparametrization

Hence, as for L_1 , we deduce that L_2 is a lens space $L(qz - ry; p - (\lambda q + \mu r)x)$ and therefore $\pi_1(\Omega_2, b) \cong \pi_1(L_2) = \langle [\tilde{G}_3 \star b] \rangle / \langle (qz - ry)[\tilde{G}_3 \star b] \rangle = \mathbf{Z}_{qz-ry}$, where $\lambda y + \mu z = 1$.

In our original coordinate system it follows that $\pi_1(\Omega_2, b) \cong \langle [G(0, \mu, -\lambda) \star b] \rangle / \langle (qz - ry)[G(0, \mu, -\lambda) \star b] \rangle$, since $A(0, \mu, -\lambda)^t = \underline{e}_2$.

Consider now $\Omega_1 \cap \Omega_2$. By the same reasoning as in [Oh2], $\pi_1(\Omega_1 \cap \Omega_2, b)$ is generated by the homotopy classes $[G_1 \star b]$, $[G_2 \star b]$ and $[G_3 \star b]$. Furthermore, $[G_1 \star b] = [G(p, q, r) \star b] = 0 \in \pi_1(\Omega_1 \cap \Omega_2, b)$.

The inclusion maps induce homomorphisms $\iota_i : \pi_1(\Omega_1 \cap \Omega_2, b) \rightarrow \pi_1(\Omega_i, b)$, $i = 1, 2$. It is clear from our previous discussion of $\pi_1(\Omega_1, b)$ that $\iota_1([G_3 \star b]) = [G_3 \star b]$ and $\iota_1([G_2 \star b]) = [G_2 \star b] = 0$. On the other hand, since A^{-1} gives a reparametrization of T^3 ,

$$\begin{aligned} \iota_2([G_3 \star b]) &= \iota_2(-y[G(0, \mu, -\lambda) \star b] + \lambda[G(x, y, z) \star b] - \mu x[G_1 \star b]) \\ &= -y\iota_2([G(0, \mu, -\lambda) \star b]) + \lambda\iota_2([G(x, y, z) \star b]) \\ &= -y[G(0, \mu, -\lambda) \star b] + \lambda[G(x, y, z) \star b] \\ &= -y[G(0, \mu, -\lambda) \star b] \end{aligned}$$

since $[G(x, y, z) \star b] = 0 \in \pi_1(\Omega_2, b)$. Similarly $\iota_2([G_2 \star b]) = z[G(0, \mu, -\lambda) \star b]$.

Let $[\alpha] = [G_3 \star b] \in \pi_1(\Omega_1, b)$ and $[\beta] = [G(0, \mu, -\lambda) \star b] \in \pi_1(\Omega_2, b)$. Then, by van Kampen's Theorem

$$\begin{aligned} \pi_1(M, b) &= (\pi_1(\Omega_1, b) * \pi_1(\Omega_2, b)) / \langle \iota_1([G_i \star b]) = \iota_2([G_i \star b]), i = 2, 3 \rangle \\ &= \langle [\alpha], [\beta] \mid r[\alpha] = 0, (qz - ry)[\beta] = 0 \rangle / \langle [\alpha] = -y[\beta], z[\beta] = 0 \rangle \\ &= \langle [\beta] \mid z[\beta] = 0, ry[\beta] = 0 \rangle \\ &= \langle [\beta] \mid \gcd(z, ry)[\beta] = 0 \rangle \\ &= \mathbf{Z}_{\gcd(r, z)}, \quad \text{since } \gcd(y, z) = 1. \end{aligned}$$

□

Corollary 4.2. *If T^3 acts smoothly and effectively on a smooth, compact, simply connected 5-manifold M^5 with (legally) weighted orbit space as in Figure 4, then*

$$(4.1) \quad \gcd(r, z) = 1, \quad \gcd(p, r) = 1, \quad \gcd(y, z) = 1 \quad \text{and} \quad \gcd(py - qx, rx - pz, qz - ry) = 1.$$

Proof. The fact that $\gcd(r, z) = 1$ follows directly from Proposition 4.1. From the discussion in Section 2 the orbit space is legally weighted if and only if there exist triples $\underline{y}_i = (\alpha_i, \beta_i, \gamma_i) \in \mathbf{Z}^3$, $i = 1, 2, 3, 4$, such that

$$\det \begin{pmatrix} \underline{y}_1 \\ \underline{x}_1 \\ \underline{x}_2 \end{pmatrix} = \pm 1, \quad \det \begin{pmatrix} \underline{y}_2 \\ \underline{x}_2 \\ \underline{x}_3 \end{pmatrix} = \pm 1, \quad \det \begin{pmatrix} \underline{y}_3 \\ \underline{x}_3 \\ \underline{x}_4 \end{pmatrix} = \pm 1 \quad \text{and} \quad \det \begin{pmatrix} \underline{y}_4 \\ \underline{x}_4 \\ \underline{x}_1 \end{pmatrix} = \pm 1.$$

This is possible if and only if the other three gcd conditions hold. □

Suppose that we have a weighted orbit space as in Figure 4 and that the conditions in (4.1) hold. Set $a = z$, $b = -(rx - pz)$, $c = r$ and $d = qz - ry$. Then $\gcd(a, c) = 1$, $\gcd(a, d) = 1$ and $\gcd(b, c) = 1$. Let $m, n \in \mathbf{Z}$ such that $am + cn = 1$.

Then $(am + cn)b = b = -cx + ap$, hence $c(x + bn) = a(p - bm)$. Therefore, since $\gcd(a, c) = 1$, $x = -bn - ak$ and $p = bm - ck$, for some $k \in \mathbf{Z}$. Similarly, it follows that $y = -dn - al$ and $q = dm - cl$, for some $\ell \in \mathbf{Z}$. Further, we find that $\gcd(b, d, py - qx) = 1$ implies that $\gcd(b, d) = 1$.

Therefore all possible legally weighted orbit spaces as in Figure 4 for a smooth, effective T^3 action on the simply connected manifolds $\mathbf{S}^3 \times \mathbf{S}^2$ and $\mathbf{S}^3 \tilde{\times} \mathbf{S}^2$ are given by the diagram




Figure 7: Possible legally weighted orbit spaces for $\mathbf{S}^3 \times \mathbf{S}^2$ and $\mathbf{S}^3 \tilde{\times} \mathbf{S}^2$

where $\gcd(a, c) = \gcd(a, d) = \gcd(b, c) = \gcd(b, d) = 1$, hence $\gcd(ab, cd) = 1$, and $k, \ell, m, n \in \mathbf{Z}$ with $am + cn = 1$.

5. FREE ACTIONS ON $\mathbf{S}^3 \times \mathbf{S}^3$

Recall that $\mathbf{S}^3 \times \mathbf{S}^3$ is a Lie group consisting of pairs of unit quaternions, i.e. pairs $(\begin{smallmatrix} q_1 \\ q_2 \end{smallmatrix})$ where $q_n = \alpha_n + \beta_n j$, with $\alpha_n, \beta_n \in \mathbf{C}$, $|\alpha_n|^2 + |\beta_n|^2 = 1$, for $n = 1, 2$. Recall that $ij = -ji$ and so, in particular, $\beta j = j \bar{\beta}$ for all $\beta \in \mathbf{C}$.

Define $\mathbf{S}_j^1 := \{\beta j \mid \beta \in \mathbf{C}, |\beta| = 1\} \subset \mathbf{S}^3$. The image under the quotient map of points $(\begin{smallmatrix} q_1 \\ q_2 \end{smallmatrix}) \in \mathbf{S}^3 \times \mathbf{S}^3$ and subgroups $H \subset \mathbf{S}^3 \times \mathbf{S}^3$ will always be denoted by $[\begin{smallmatrix} q_1 \\ q_2 \end{smallmatrix}]$ and $[H]$ respectively.

Let \langle, \rangle_0 be a bi-invariant metric on $\mathbf{S}^3 \times \mathbf{S}^3$. The usual isometric T^4 action on $(\mathbf{S}^3 \times \mathbf{S}^3, \langle, \rangle_0)$, namely

$$(u, v, w, z) \star \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \begin{pmatrix} u q_1 \bar{v} \\ w q_2 \bar{z} \end{pmatrix}, \quad (u, v, w, z) \in T^4, \quad q_1, q_2 \in \mathbf{S}^3,$$

is not effective. Consider instead the T^4 action given by

$$(5.1) \quad (u, v, w, z) \star \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \begin{pmatrix} u \alpha_1 + v \beta_1 j \\ w \alpha_2 + z \beta_2 j \end{pmatrix}, \quad (u, v, w, z) \in T^4, \quad q_1, q_2 \in \mathbf{S}^3.$$

This action is clearly effective. Moreover, it is an isometric action since it may be rewritten as a (well-defined) two-sided action as follows:

$$(u, v, w, z) \star \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \begin{pmatrix} u^{\frac{1}{2}} v^{\frac{1}{2}} q_1 u^{\frac{1}{2}} \bar{v}^{\frac{1}{2}} \\ w^{\frac{1}{2}} z^{\frac{1}{2}} q_2 w^{\frac{1}{2}} \bar{z}^{\frac{1}{2}} \end{pmatrix}.$$

Remark 5.1. Define $S(U(2) \times U(2)) := \{(A, B) \in U(2) \times U(2) \mid \det(A) = \det(B)\}$. Then the action (5.1) can, in fact, be thought of as an action of $T^4 \subset S(U(2) \times U(2))^2$ on $SU(2) \times SU(2)$ since, for example, an \mathbf{S}^1 action $z \star q = z^{\frac{2k+1}{2}} q \bar{z}^{\frac{2\ell+1}{2}}$, $q = \alpha + \beta j \in \mathbf{S}^3$, $z \in \mathbf{S}^1$, is equivalent to an action of $\mathbf{S}^1 \subset S(U(2) \times U(2))$ on $SU(2)$ via

$$z \star \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} = \begin{pmatrix} z^k & \\ & \bar{z}^{k+1} \end{pmatrix} \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} \begin{pmatrix} \bar{z}^\ell & \\ & z^{\ell+1} \end{pmatrix}.$$

Consider the tuples $(a, b, c, d), (-n, k, m, \ell) \in \mathbf{Z}^4$, such that $am + cn = 1$. As

$$\det \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ -n & k & m & \ell \\ a & b & c & d \end{pmatrix} = am + cn = 1$$

we may reparametrize the effective, isometric action (5.1) via

$$(u, v, w, z) \mapsto (z^a \bar{w}^n, uz^b w^k, z^c w^m, vz^d w^\ell)$$

to give an effective, isometric T^4 action on $(\mathbf{S}^3 \times \mathbf{S}^3, \langle, \rangle_0)$ via

$$(5.2) \quad (u, v, w, z) \star \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \begin{pmatrix} z^a \bar{w}^n \alpha_1 + uz^b w^k \beta_1 j \\ z^c w^m \alpha_2 + vz^d w^\ell \beta_2 j \end{pmatrix}.$$

In particular, if the circle (resp. torus) given by the z -coordinate (resp. (w, z) -coordinates) acts freely on $\mathbf{S}^3 \times \mathbf{S}^3$, then there is an induced effective, isometric action on the quotient by the (u, v, w) -torus (resp. (u, v) -torus).

This raises the question of when the z -circle and (w, z) -torus act freely. It is easy to check that \mathbf{S}^1 acts freely on $\mathbf{S}^3 \times \mathbf{S}^3$ via

$$(5.3) \quad z \star \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \begin{pmatrix} z^a \alpha_1 + z^b \beta_1 j \\ z^c \alpha_2 + z^d \beta_2 j \end{pmatrix}$$

if and only if $\gcd(a, c) = \gcd(a, d) = \gcd(b, c) = \gcd(b, d) = 1$, that is, if and only if $\gcd(ab, cd) = 1$. We denote the quotient $(\mathbf{S}^3 \times \mathbf{S}^3, \langle, \rangle_0) // \mathbf{S}^1$ by $M_{a,b,c,d}^5$.

DeVito [DV] has classified up to diffeomorphism all possible biquotients $(\mathbf{S}^3 \times \mathbf{S}^3) // \mathbf{S}^1$. More precisely, he has shown that only $\mathbf{S}^3 \times \mathbf{S}^2$ and $\mathbf{S}^3 \tilde{\times} \mathbf{S}^2$ can arise. Since both of these manifolds satisfy $H_2(M^5; \mathbf{Z}) = \mathbf{Z}$, this classification amounts to computing the second Stiefel-Whitney class $w_2((\mathbf{S}^3 \times \mathbf{S}^3) // \mathbf{S}^1) \in H^2(M^5; \mathbf{Z}_2)$ (cf. [Ba]). A consequence of DeVito's classification is that the quotient resulting from the explicit displayed action suggested by Pavlov [Pav] and claimed to be $\mathbf{S}^3 \tilde{\times} \mathbf{S}^2$, in fact has $w_2 = 0$, hence must be diffeomorphic to $\mathbf{S}^3 \times \mathbf{S}^2$. In the paragraph preceeding this displayed action, Pavlov describes in words an action which gives the correct quotient.

It can be shown that $w_2(M_{a,b,c,d}^5) = a + b + c + d \in \mathbf{Z}_2$. Hence, together with the freeness condition, it follows that $M_{a,b,c,d}^5$ is diffeomorphic to $\mathbf{S}^3 \tilde{\times} \mathbf{S}^2$ if and only if exactly one of a, b, c or d is even (and is diffeomorphic to $\mathbf{S}^3 \times \mathbf{S}^2$ otherwise). It should be noted that although we have parametrized our \mathbf{S}^1 actions on $\mathbf{S}^3 \times \mathbf{S}^3$ differently to those in [DV], the computation of the Stiefel-Whitney class w_2 is completely analogous to the computation carried out therein, and hence has been omitted.

On the other hand, T^2 acts freely on $\mathbf{S}^3 \times \mathbf{S}^3$ via

$$(5.4) \quad (w, z) \star \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \begin{pmatrix} z^a \bar{w}^n \alpha_1 + z^b w^k \beta_1 j \\ z^c w^m \alpha_2 + z^d w^\ell \beta_2 j \end{pmatrix}$$

if and only if (without loss of generality)

$$(5.5) \quad am + cn = 1, \quad al + dn = \varepsilon_2, \quad bm - ck = \varepsilon_3 \quad \text{and} \quad bl - dk = \varepsilon_4,$$

where $\varepsilon_2, \varepsilon_3, \varepsilon_4 = \pm 1$.

Lemma 5.2. *Suppose $a, c \in \mathbf{Z}$ are relatively prime and let $m_0, n_0 \in \mathbf{Z}$ such that $am_0 + cn_0 = 1$. Then $m, n \in \mathbf{Z}$ satisfy $am + cn = 1$ if and only if there exists $x \in \mathbf{Z}$ such that $m = m_0 - cx$ and $n = n_0 + ax$.*

Proof. As the cases $a = 0$ and $c = 0$ are trivial, assume that $a, c \neq 0$. By subtracting $am + cn = 1$ from $am_0 + cn_0 = 1$ it follows that $a(m_0 - m) = -c(n_0 - n)$. But a and c are assumed to be relatively prime, hence $n_0 - n = -ax$ and $m_0 - m = cx$ for some $x \in \mathbb{Z}$. \square

Using this lemma, one can describe all possible free T^2 actions on $S^3 \times S^3$. In the case where $\varepsilon_2 \varepsilon_3 \varepsilon_4 = 1$ it turns out that, up to reparametrization of the action and diffeomorphisms of $S^3 \times S^3$ (namely $q_1 \leftrightarrow q_2$ and $q_i \mapsto \bar{q}_i$), the action has the form

$$(5.6) \quad (w, z) \star \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \begin{pmatrix} z\bar{w}^r \alpha_1 + zw^{r+\lambda} \beta_1 j \\ w\alpha_2 + w\beta_2 j \end{pmatrix}$$

where $r \in \mathbb{Z}$ and $\lambda \in \{0, 1\}$. Furthermore, $M_{r,\lambda}^4 := (S^3 \times S^3, \langle, \rangle_0) // T^2$ is diffeomorphic to $S^2 \times S^2$ for $\lambda = 0$, and to $\mathbb{CP}^2 \# -\mathbb{CP}^2$ for $\lambda = 1$. This follows from work in [DV] via a change of parameters (cf. also [Ch], [To]).

In the case where $\varepsilon_2 \varepsilon_3 \varepsilon_4 = -1$ it can be shown that, up to reparametrization of the action and diffeomorphisms of $S^3 \times S^3$, the action has the form

$$(5.7) \quad (w, z) \star \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \begin{pmatrix} z\alpha_1 + w\beta_1 j \\ \bar{z}w\alpha_2 + zw\beta_2 j \end{pmatrix}.$$

The quotient $M^4 = (S^3 \times S^3, \langle, \rangle_0) // T^2$ is diffeomorphic to $\mathbb{CP}^2 \# \mathbb{CP}^2$ (cf. [To], [DV]).

Remark 5.3. Recall that the only smooth, compact, simply connected 4-manifolds admitting both non-negative curvature and an isometric circle action are S^4 , \mathbb{CP}^2 , $\mathbb{CP}^2 \# \pm \mathbb{CP}^2$ and $S^2 \times S^2$ (cf. [KI], [SY]). The long exact homotopy sequence for a fibration $T^2 \rightarrow S^3 \times S^3 \rightarrow M^4$ shows that S^4 and \mathbb{CP}^2 cannot arise as biquotients $(S^3 \times S^3, \langle, \rangle_0) // T^2$.

6. TORUS ACTIONS ON FOUR-DIMENSIONAL BIQUOTIENTS

In the T^4 action described by (5.2), let T_{uv}^2 and T_{wz}^2 denote the 2-tori given by the (u, v) and (w, z) coordinates respectively. Consider the induced effective, isometric T_{uv}^2 action on any $M^4 = (S^3 \times S^3) // T_{wz}^2$, where T_{wz}^2 acts freely (as in (5.6) or (5.7)), namely

$$(6.1) \quad (u, v) \star \begin{bmatrix} q_1 \\ q_2 \end{bmatrix} = \begin{bmatrix} \alpha_1 + u\beta_1 j \\ \alpha_2 + v\beta_2 j \end{bmatrix}.$$

It is clear that the quotient of $S^1 \times S^1 \subset S^3 \times S^3$ under T_{wz}^2 is a point, whereas $S^1 \times S^3 \subset S^3 \times S^3$ has quotient diffeomorphic to S^2 . Similarly, $S^1 \times S_j^1$, $S_j^1 \times S^1$ and $S_j^1 \times S_j^1$ quotient to points, while $S_j^1 \times S^3$, $S^3 \times S^1$ and $S^3 \times S_j^1$ have quotients diffeomorphic to S^2 .

For each of the biquotients $(S^3 \times S^3, \langle, \rangle_0) // T_{wz}^2$ we will determine the fixed-point set of the corresponding effective, isometric T_{uv}^2 action, as well as any additional isotropy that may arise. As mentioned in Section 2, the only possible isotropy groups of an effective T^2 action on a four-dimensional manifold are S^1 and T^2 .

Lemma 6.1. *Let T_{wz}^2 act on $(S^3 \times S^3, \langle, \rangle_0)$ via (5.6) and let T_{uv}^2 act on $M_{r,\lambda}^4 = (S^3 \times S^3, \langle, \rangle_0) // T_{wz}^2$ via (6.1). Then the fixed points of the action are the four points $[S^1 \times S^1]$, $[S_j^1 \times S^1]$, $[S^1 \times S_j^1]$, $[S_j^1 \times S_j^1] \in S^3 \times S^3 // T_{k,\varepsilon}^2$, while a point has S^1 isotropy if and only if it lies in one of the four two-spheres $[S^1 \times S^3]$, $[S_j^1 \times S^3]$, $[S^3 \times S^1]$ or $[S^3 \times S_j^1]$. In particular, the*

S^1 isotropy subgroups of T_{uv}^2 are arranged according to the diagram

$$(6.2) \quad \begin{array}{ccccc} & & \{(u,1)\} & & \\ & & \text{---} & & \\ & & [S^1 \times S^3] & & \\ & & \text{---} & & \\ [S^1 \times S^1] & \cdots & [S^1 \times S_j^1] & \cdots & [S^1 \times S_j^1] \\ \vdots & & \vdots & & \vdots \\ \{(1,v)\} & [S^3 \times S^1] & & [S^3 \times S_j^1] & \{(v^{2r+\lambda}, v)\} \\ \vdots & & \vdots & & \vdots \\ [S_j^1 \times S^1] & \cdots & [S_j^1 \times S^3] & \cdots & [S_j^1 \times S_j^1] \\ & & \{(u,1)\} & & \end{array}$$

Proof. Suppose that $\begin{bmatrix} q_1 \\ q_2 \end{bmatrix} \in M_{r,\lambda}^4$ is fixed by some element of T_{uv}^2 . That is, there exist $(w, z) \in T_{wz}^2$ such that

$$\begin{pmatrix} \alpha_1 + \beta_1 j \\ \alpha_2 + \beta_2 j \end{pmatrix} = \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \begin{pmatrix} z\bar{w}^r \alpha_1 + uz\bar{w}^{r+\lambda} \beta_1 j \\ w\alpha_2 + wv\beta_2 j \end{pmatrix}.$$

Then each of the following four conditions must hold:

- $\alpha_1 = 0$ or $z\bar{w}^r = 1$;
- $\beta_1 = 0$ or $uz\bar{w}^{r+\lambda} = 1$;
- $\alpha_2 = 0$ or $w = 1$;
- $\beta_2 = 0$ or $wv = 1$.

First suppose that $\alpha_1 = \alpha_2 = 0$. Then $w = \bar{v}$ and $1 = uz\bar{w}^{r+\lambda} = uz\bar{v}^{r+\lambda}$, since $|\beta_1| = |\beta_2| = 1$. Therefore $z = \bar{u}v^{r+\lambda}$ and hence for all (u, v) we have $(u, v) \star [S_j^1 \times S_j^1] = [S_j^1 \times S_j^1]$. That is, $[S_j^1 \times S_j^1]$ is a fixed point of the T_{uv}^2 action.

The analogous computations in the cases $\beta_1 = \alpha_2 = 0$, $\alpha_1 = \beta_2 = 0$ and $\beta_1 = \beta_2 = 0$ show that $[S^1 \times S_j^1]$, $[S_j^1 \times S^1]$ and $[S^1 \times S^1]$, respectively, are also fixed points of the T_{uv}^2 action.

By effectiveness of the T_{uv}^2 action, it is clear that whenever all of $\alpha_1, \alpha_2, \beta_1$ and β_2 are non-zero, the isotropy group is trivial.

Now suppose that $\alpha_2 = 0$ and $\alpha_1, \beta_1 \neq 0$. Then $w = \bar{v}$ and $z\bar{w}^r = 1 = uz\bar{w}^{r+\lambda}$. Hence $z = \bar{v}^r = \bar{u}v^{r+\lambda}$, from which it follows that $u = v^{2r+\lambda}$. Therefore each point of $[S^3 \times S_j^1]$ is fixed by the circle $\{(v^{2r+\lambda}, v)\}$ in T_{uv}^2 .

Similarly we find that the points of $[S^1 \times S^3]$, $[S_j^1 \times S^3]$ and $[S^3 \times S^1]$ are fixed by the circles $\{(u, 1)\}$, $\{(u, 1)\}$ and $\{(1, v)\}$ in T_{uv}^2 respectively. \square

Proposition 6.2. *Every smooth, effective T^2 action on $S^2 \times S^2$, $\mathbb{CP}^2 \# -\mathbb{CP}^2$ or $\mathbb{CP}^2 \# \mathbb{CP}^2$ is equivariantly diffeomorphic to an effective, isometric T_{uv}^2 action on the corresponding biquotient $(S^3 \times S^3, \langle \cdot, \cdot \rangle_0) // T_{wz}^2$.*

Proof. From the discussion in Section 3 we know that there is a unique smooth, effective T^2 action on $\mathbb{CP}^2 \# \mathbb{CP}^2$ up to equivariant diffeomorphism. It is clear, therefore, that this action must correspond to the effective, isometric T_{uv}^2 action on the biquotient $\mathbb{CP}^2 \# \mathbb{CP}^2 = S^3 \times S^3 // T_{wz}^2$, for T_{wz}^2 acting via (5.7).

From diagram (6.2) in Lemma 6.1 it follows that the weighted orbit space of a T_{uv}^2 action on $M_{r,\lambda}^4$ (that is, the diagram of slopes of S^1 isotropy groups in T_{uv}^2) is given by

$$(6.3) \quad \begin{array}{ccc} [S^1 \times S^1] & \xrightarrow{(1,0)} & [S^1 \times S_j^1] \\ \vdots (0,1) & & \vdots (2r+\lambda,1) \\ [S_j^1 \times S^1] & \xrightarrow{(1,0)} & [S_j^1 \times S_j^1] \end{array}$$

where $r \in \mathbf{Z}$ and $\lambda \in \{0, 1\}$. When $\lambda = 0$ (resp. $\lambda = 1$) it is clear from the discussion in Section 3 that all possible weighted orbit spaces for $S^2 \times S^2$ (resp. $\mathbf{CP}^2 \# -\mathbf{CP}^2$) are achieved. \square

Remark 6.3. A simple observation is that for each pair $(r, s) \in \mathbf{Z}^2$ there is a three-dimensional orbifold $X_{r,s}^3$ such that $S^2 \times S^2$ and $\mathbf{CP}^2 \# -\mathbf{CP}^2$ arise as the total spaces of particular S^1 -bundles over $X_{r,s}^3$. Indeed, the three-torus action on $(S^3 \times S^3, \langle, \rangle_0)$ defined by

$$(u, w, z) \star \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \begin{pmatrix} z\bar{w}^r \bar{u}^s \alpha_1 + zw^{r+1} u^s \beta_1 j \\ w\bar{u} \alpha_2 + wu \beta_2 j \end{pmatrix}$$

is effective and isometric. Denote the quotient space $(S^3 \times S^3, \langle, \rangle_0) / T_{uwz}^3$ by $X_{r,s}^3$. It follows from the work of Perelman [Pe1, Pe2] that the orbifold $X_{r,s}^3$ is homeomorphic to S^3 as a topological manifold. The subtori given by restriction to the (z, w) and (z, u) coordinates are both of the form (5.6) and have quotients $\mathbf{CP}^2 \# -\mathbf{CP}^2$ and $S^2 \times S^2$ respectively. In each case the remaining effective and isometric $(u$ or $w)$ circle action on the 4-manifold has isotropy subgroups arranged according to the diagram

$$\begin{array}{ccccc} [S^1 \times S^1] & \xrightarrow[\mathbf{Z}_2]{[S^1 \times S^3]} & [S^1 \times S_j^1] \\ \vdots \mathbf{Z}_{|2(r+s)+1|} & & \vdots \mathbf{Z}_{|2(r-s)+1|} \\ [S^3 \times S^1] & & [S^3 \times S_j^1] \\ \vdots & & \vdots \\ [S_j^1 \times S^1] & \xrightarrow[\mathbf{Z}_2]{[S_j^1 \times S^3]} & [S_j^1 \times S_j^1] \end{array}$$

where the vertices are fixed points of the action. In the context of Fintushel's classification of smooth, effective circle actions on smooth, compact, simply-connected 4-manifolds via Seifert invariants (α_i, β_i) [Fi1], this shows that there are infinitely many smooth, effective circle actions on $\mathbf{CP}^2 \# -\mathbf{CP}^2$ and $S^2 \times S^2$ having the same α_i invariants (i.e. the orders of the isotropy groups), and so these actions must be distinguished by their (harder to compute) β_i invariants.

Remark 6.4. In the cases of $\mathbf{CP}^2 \# \pm \mathbf{CP}^2$ and $S^2 \times S^2$, Corollary B can also be proven directly, without appealing to Theorem A, by determining (up to equivariant diffeomorphism) all possible isometric circle actions on these manifolds in terms of Fintushel's classification of smooth, effective circle actions on smooth, compact, simply connected 4-manifolds [Fi1] and then performing computations similar to those in this section to

show that each of these actions is realised by an effective, isometric circle action on the corresponding normal biquotient.

For fixed point homogeneous circle actions, that is, those where the fixed point set of the action has a two-dimensional component (cf. [GS2] and [GG]), all possible weighted orbit spaces (hence all such actions on these manifolds) were determined in [GG]. One can then reach the conclusion of Corollary B without using the work of Grove and Wilking [GW]. To determine the weighted orbit spaces for circle actions which are not fixed point homogeneous, one must combine Fintushel's classification with Theorem 2.4 in [GW].

7. TORUS ACTIONS ON FIVE-DIMENSIONAL BIQUOTIENTS

From Oh's classification of smooth cohomogeneity-two torus actions on smooth 5-manifolds [Oh2], we know that there is, up to equivariant diffeomorphism, a unique smooth T^3 action on S^5 . This is, of course, realized by the linear (isometric) T^3 action on the homogeneous space $S^5 = SO(6)/SO(5)$. As we are interested only in those simply connected 5-manifolds which admit a T^3 -invariant metric with non-negative curvature, that is, S^5 , $S^3 \times S^2$ and $S^3 \tilde{\times} S^2$ (cf. [GGS]), we may restrict our attention from now on to smooth T^3 actions on the manifolds $S^3 \times S^2$ and $S^3 \tilde{\times} S^2$.

Let S^1 act freely and isometrically on $(S^3 \times S^3, \langle \cdot, \cdot \rangle_0)$ via (5.3). Denote the quotient $(S^3 \times S^3, \langle \cdot, \cdot \rangle_0)/S^1$ by $M_{a,b,c,d}^5$. By the discussion in Section 5, $M_{a,b,c,d}^5$ is diffeomorphic to $S^3 \times S^2$ or $S^3 \tilde{\times} S^2$ depending on the parity of $a + b + c + d$.

Consider the effective, isometric T^4 action on $(S^3 \times S^3, \langle \cdot, \cdot \rangle_0)$ in (5.2). The complementary three-dimensional torus T_{uvw}^3 to the z -circle in T^4 then acts on $M_{a,b,c,d}^5$ effectively and isometrically via

$$(7.1) \quad (u, v, w) \star \begin{bmatrix} q_1 \\ q_2 \end{bmatrix} = \begin{bmatrix} \bar{w}^n \alpha_1 + u w^k \beta_1 j \\ w^m \alpha_2 + v w^\ell \beta_2 j \end{bmatrix}$$

where $(u, v, w) \in T_{uvw}^3$, $[q_1] \in M_{a,b,c,d}^5$, $k, \ell, m, n \in \mathbf{Z}$ and $am + cn = 1$.

Lemma 7.1. *Let T_{uvw}^3 act effectively and isometrically on $M_{a,b,c,d}^5$ via the action (7.1). Then the four points $[S^1 \times S^1]$, $[S_j^1 \times S^1]$, $[S^1 \times S_j^1]$, $[S_j^1 \times S_j^1] \in M_{a,b,c,d}^5$ each have T^2 isotropy, while a point has S^1 isotropy if and only if it lies in one of $[S^1 \times S^3]$, $[S_j^1 \times S^3]$, $[S^3 \times S^1]$ or $[S^3 \times S_j^1]$. In particular, the S^1 isotropy subgroups of T_{uvw}^3 are arranged according to the diagram:*

$$(7.2) \quad \begin{array}{ccc} [S^1 \times S^1] & \xrightarrow[\{S^1 \times S^3\}]{\{(1,v,1)\}} & [S^1 \times S_j^1] \\ \vdots & & \vdots \\ \{(u,1,1)\} [S^3 \times S^1] & & [S^3 \times S_j^1] \{(\bar{w}^{bn+ak}, \bar{w}^{dn+c\ell}, w^a)\} \\ \vdots & & \vdots \\ [S_j^1 \times S^1] & \xrightarrow[\{(w^{bm-ck}, w^{dm-c\ell}, w^c)\}]{[S_j^1 \times S^3]} & [S_j^1 \times S_j^1] \end{array}$$

Proof. Suppose that $[q_1]$ is fixed by some element (u, v, w) of T_{uvw}^3 . That is, there exists $z \in S^1$ such that

$$\begin{pmatrix} \alpha_1 + \beta_1 j \\ \alpha_2 + \beta_2 j \end{pmatrix} = \begin{pmatrix} z^a \bar{w}^n \alpha_1 + u z^b w^k \beta_1 j \\ z^c w^m \alpha_2 + v z^d w^\ell \beta_2 j \end{pmatrix}.$$

Then each of the following four conditions must hold:

- $\alpha_1 = 0$ or $z^a \bar{w}^n = 1$;
- $\beta_1 = 0$ or $uz^b \bar{w}^k = 1$;
- $\alpha_2 = 0$ or $z^c w^m = 1$;
- $\beta_2 = 0$ or $vz^d \bar{w}^\ell = 1$.

By effectiveness of the T_{uvw}^3 action, it is clear that whenever all of $\alpha_1, \alpha_2, \beta_1$ and β_2 are non-zero, the isotropy group is trivial.

Suppose first that $\beta_1 = \beta_2 = 0$. Then $z^a \bar{w}^n = 1$ and $z^c w^m = 1$, from which it follows that $w = z = 1$, since $am + cn = 1$. Thus the isotropy subgroup of the point $[\mathbf{S}^1 \times \mathbf{S}^1]$ is given by

$$T^2 \cong \{(u, v, 1)\} \subset T_{uvw}^3.$$

If $\alpha_1 = \alpha_2 = 0$, then $u = \bar{z}^b \bar{w}^k$ and $v = \bar{z}^d \bar{w}^\ell$, since $|\beta_1| = |\beta_2| = 1$. Therefore the isotropy subgroup of the point $[\mathbf{S}_j^1 \times \mathbf{S}_j^1]$ is

$$T^2 \cong \{(\bar{z}^b \bar{w}^k, \bar{z}^d \bar{w}^\ell, w)\} \subset T_{uvw}^3.$$

Suppose now that $\beta_1 = \alpha_2 = 0$. Then $z^a \bar{w}^n = 1$ and $v = \bar{z}^d \bar{w}^\ell$. Let $\zeta \in \mathbf{S}^1$ such that $\zeta^a = w$. It follows by setting $z = \zeta^n$ that $z^a = w^n$ and $v = \bar{\zeta}^{dn+a\ell}$. Hence the isotropy subgroup of the point $[\mathbf{S}^1 \times \mathbf{S}_j^1]$ is given by

$$T^2 \cong \{(u, \bar{\zeta}^{dn+a\ell}, \zeta^a)\} \subset T_{uvw}^3.$$

We remark that $\gcd(a, dn + a\ell) = 1$, since $\gcd(ab, cd) = 1$ and $am + cn = 1$.

The analogous computation for $\alpha_1 = \beta_2 = 0$ shows that the isotropy subgroup of the point $[\mathbf{S}_j^1 \times \mathbf{S}^1]$ is given by

$$T^2 \cong \{(\zeta^{bm-ck}, v, \zeta^c)\} \subset T_{uvw}^3.$$

Suppose, on the other hand, that $\beta_1 = 0$ and $\alpha_2, \beta_2 \neq 0$. Then $z^a \bar{w}^n = 1, z^c w^m = 1$ and $v = \bar{z}^d \bar{w}^\ell$. As before it follows from $am + cn = 1$ that $w = z = 1$, hence $v = 1$. Therefore the isotropy subgroup of a generic point in $[\mathbf{S}^1 \times \mathbf{S}^3]$ is

$$\mathbf{S}^1 \cong \{(u, 1, 1)\} \subset T_{uvw}^3.$$

Similarly, generic points in $[\mathbf{S}^3 \times \mathbf{S}^1]$ (i.e. points for which $\beta_2 = 0$ and $\alpha_1, \beta_1 \neq 0$) have isotropy subgroup

$$\mathbf{S}^1 \cong \{(1, v, 1)\} \subset T_{uvw}^3.$$

If now $\alpha_2 = 0$ and $\alpha_1, \beta_1 \neq 0$, then $v = \bar{z}^d \bar{w}^\ell, u = \bar{z}^b \bar{w}^k$ and $z^a \bar{w}^n = 1$. As above, let $\zeta \in \mathbf{S}^1$ such that $\zeta^a = w$ and set $z = \zeta^n$. Then $u = \bar{\zeta}^{bn+ak}$ and $v = \bar{\zeta}^{dn+a\ell}$. Therefore a generic point in $[\mathbf{S}^3 \times \mathbf{S}_j^1]$ has isotropy subgroup

$$\mathbf{S}^1 \cong \{(\bar{\zeta}^{bn+ak}, \bar{\zeta}^{dn+a\ell}, \zeta^a)\} \subset T_{uvw}^3.$$

The computation to show that a generic point in $[\mathbf{S}_j^1 \times \mathbf{S}^3]$ (i.e. $\alpha_1 = 0$ and $\alpha_2, \beta_2 \neq 0$) has isotropy subgroup

$$\mathbf{S}^1 \cong \{(\zeta^{bm-ck}, \zeta^{dm-cl}, \zeta^c)\} \subset T_{uvw}^3$$

is completely analogous. \square

Proposition 7.2. *Every smooth, effective T^3 action on either $\mathbf{S}^3 \times \mathbf{S}^2$ or $\mathbf{S}^3 \tilde{\times} \mathbf{S}^2$ is equivariantly diffeomorphic to an effective, isometric T_{uvw}^3 action on the corresponding biquotient $M_{a,b,c,d}^5 = (\mathbf{S}^3 \times \mathbf{S}^3, \langle \cdot, \cdot \rangle_0) // \mathbf{S}^1$.*

Proof. From diagram 7.2 in Lemma 7.1 it follows that the weighted orbit space of an effective, isometric T^3 action on $M_{a,b,c,d}^5$ (that is, the diagram of slopes of S^1 isotropy groups in T_{uvw}^3) is given by

$$(7.3) \quad \begin{array}{ccc} [S^1 \times S^1] & \xrightarrow{(1,0,0)} & [S^1 \times S_j^1] \\ \vdots (0,1,0) & & \vdots (-bn-ak, -dn-a\ell, a) \\ [S_j^1 \times S^1] & \xrightarrow{(bm-ck, dm-cl, c)} & [S_j^1 \times S_j^1] \end{array}$$

where $\gcd(ab, cd) = 1$, $am + cn = 1$, and $k, \ell \in \mathbf{Z}$. It is then clear from the discussion in Section 4 that all possible (legally) weighted orbit spaces for $S^3 \times S^2$ and $S^3 \tilde{\times} S^2$ are achieved. \square

8. PRINCIPAL CIRCLE BUNDLES

Let B be an arbitrary simply connected manifold. It is well known that oriented (hence principal) circle bundles over B are classified by their Euler classes in the second integral cohomology $H^2(B; \mathbf{Z})$. Since B is simply connected, the total space P of the bundle is simply connected if and only if the Euler characteristic is primitive, i.e. a generator of $H^2(B; \mathbf{Z})$.

As a consequence of their analysis of oriented circle bundles over compact, simply connected 4-manifolds, Duan and Liang [DL] have shown that if P is simply connected and the total space of a principal circle bundle over $S^2 \times S^2$ or $\mathbf{CP}^2 \# \pm \mathbf{CP}^2$, then it must be diffeomorphic to one of $S^3 \times S^2$ or $S^3 \tilde{\times} S^2$. On the other hand, Grove and Ziller [GZ, Theorem 4.5] have observed that simply connected principal circle bundles over $S^2 \times S^2$ and $\mathbf{CP}^2 \# \pm \mathbf{CP}^2$ all arise by considering circle sub-actions of free (isometric) torus actions on $S^3 \times S^3$, namely

$$S^1 = T^2 / S^1 \longrightarrow (S^3 \times S^3) / S^1 \longrightarrow (S^3 \times S^3) // T^2.$$

From the discussions in previous sections, one can make the following observation.

Proposition 8.1. *Suppose T^2 acts freely and isometrically on $S^3 \times S^3$. Embed S^1 into T^2 via $z \mapsto (z^p, z^q)$, $p, q \in \mathbf{Z}$, $\gcd(p, q) = 1$, and denote this circle subgroup $S_{p,q}^1$. Consider the principal S^1 -bundle*

$$S^1 = T^2 / S_{p,q}^1 \longrightarrow X_{p,q}^5 := (S^3 \times S^3) / S_{p,q}^1 \longrightarrow M^4 := (S^3 \times S^3) // T^2.$$

- (a) *If $M^4 = S^2 \times S^2$, then $X_{p,q}^5 = S^3 \times S^2$ for all p, q .*
- (b) *If $M^4 = \mathbf{CP}^2 \# -\mathbf{CP}^2$, then $X_{p,q}^5 = S^3 \times S^2$ for p even and $X_{p,q}^5 = S^3 \tilde{\times} S^2$ for p odd.*
- (c) *If $M^4 = \mathbf{CP}^2 \# \mathbf{CP}^2$, then $X_{p,q}^5 = S^3 \times S^2$ for $p + q$ even and $X_{p,q}^5 = S^3 \tilde{\times} S^2$ for $p + q$ odd.*

Proof. Recall that every free, isometric T^2 action on $S^3 \times S^3$ is equivalent to one of the T_{wz}^2 actions (5.6) or (5.7) described in Section 3. First consider $S_{p,q}^1$ as a sub-action of (5.6). Then $S_{p,q}^1$ acts on $S^3 \times S^3$ via

$$z \star \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \begin{pmatrix} z^{q-pr} \alpha_1 + z^{q+p(r+\lambda)} \beta_1 j \\ z^p \alpha_2 + z^p \beta_2 j \end{pmatrix}.$$

From (5.3) it follows that $a = q - pr$, $b = q + p(r + \lambda)$, $c = p$ and $d = p$. Since $w_2(X_{p,q}^5) = a + b + c + d \in \mathbf{Z}_2$, in the cases of $M^4 = \mathbf{S}^2 \times \mathbf{S}^2$ and $M^4 = \mathbf{CP}^2 \# -\mathbf{CP}^2$ we are done. The case $M^4 = \mathbf{CP}^2 \# \mathbf{CP}^2$ is similar. \square

It is now natural to ask whether every free (isometric) \mathbf{S}^1 action on $\mathbf{S}^3 \times \mathbf{S}^3$ extends to a free (isometric) \mathbf{T}^2 action, and hence defines a principal circle bundle as above. This amounts to asking whether the effective \mathbf{T}^3 action in (7.1) contains a free circle sub-action.

Lemma 8.2. *Let $M_{a,b,c,d}^5 = (\mathbf{S}^3 \times \mathbf{S}^3, \langle, \rangle_0) // \mathbf{S}^1$ be as in Section 5 and let \mathbf{T}_{uvw}^3 act on $M_{a,b,c,d}^5$ via the action (7.1). Up to reparametrization, a circle embedded in \mathbf{T}_{uvw}^3 via $u \mapsto (u^p, u^q, u^r)$, $\gcd(p, q, r) = 1$, acts freely on $M_{a,b,c,d}^5$ if and only if $r = 1$ and*

$$a(q + \ell) + dn = \pm 1, \quad c(p + k) - bm = \pm 1, \quad \text{and} \quad b(q + \ell) - d(p + k) = \pm 1$$

for some choice of signs. Furthermore, a necessary condition for such an \mathbf{S}^1 action to be free is

$$(8.1) \quad bd \pm ac \pm ad \pm bc = 0$$

for some choice of signs.

Proof. The \mathbf{S}^1 action on $M_{a,b,c,d}^5$ is given by

$$u \star \begin{bmatrix} q_1 \\ q_2 \end{bmatrix} = \begin{bmatrix} \bar{u}^{rn} \alpha_1 + u^{p+rk} \beta_1 j \\ u^{rm} \alpha_2 + u^{q+r\ell} \beta_2 j \end{bmatrix}$$

We may assume that each of $q_n \in \mathbf{S}^3$, $n = 1, 2$, is in one of \mathbf{S}^1 or \mathbf{S}_j^1 , since allowing both α_n and β_n to be non-zero will only increase the number of freeness conditions to be satisfied. Then $u \star \begin{bmatrix} q_1 \\ q_2 \end{bmatrix} = \begin{bmatrix} q_1 \\ q_2 \end{bmatrix}$ if and only if there exists some $z \in \mathbf{S}^1$ such that

$$\begin{pmatrix} \alpha_1 + \beta_1 j \\ \alpha_2 + \beta_2 j \end{pmatrix} = \begin{pmatrix} \bar{u}^{rn} z^a \alpha_1 + u^{p+rk} z^b \beta_1 j \\ u^{rm} z^c \alpha_2 + u^{q+r\ell} z^d \beta_2 j \end{pmatrix}.$$

That is, if and only if there is some $z \in \mathbf{S}^1$ such that

- (i) $\bar{u}^{rn} z^a = 1$ and $u^{rm} z^c = 1$; or
- (ii) $\bar{u}^{rn} z^a = 1$ and $u^{q+r\ell} z^d = 1$; or
- (iii) $u^{p+rk} z^b = 1$ and $u^{rm} z^c = 1$; or
- (iv) $u^{p+rk} z^b = 1$ and $u^{q+r\ell} z^d = 1$.

In each case, if $u = 1$ then $z = 1$ by freeness of the z action. By (i) we have $\bar{u}^{rcn} = \bar{z}^{ac} = u^{ram}$, from which it follows that $1 = u^{r(am+cn)} = u^r$. Hence a necessary condition for freeness is that $r = \pm 1$. Without loss of generality, assume that $r = 1$.

Similarly, (ii) implies that $\bar{u}^{dn} = \bar{z}^{ad} = u^{a(q+\ell)}$, hence $u^{dn+a(q+\ell)} = 1$. Then $u = 1$ if and only if $dn + a(q + \ell) = \pm 1$. The conditions (iii) and (iv) yield $c(p + k) - bm = \pm 1$ and $b(q + \ell) - d(p + k) = \pm 1$ in the same way.

Finally, equation (8.1) arises by noticing that (for an appropriate choice of signs)

$$\begin{aligned} bd \pm ac \pm ad \pm bc &= bd(am + cn) + ac(b(q + \ell) - d(p + k)) \\ &\quad + ad(c(p + k) - bm) - bc(dn + a(q + \ell)) \\ &= 0. \end{aligned}$$

\square

Corollary 8.3. *There exist infinitely many free (isometric) \mathbf{S}^1 actions on $\mathbf{S}^3 \times \mathbf{S}^3$ which do not extend to a free (isometric) \mathbf{T}^2 action, and infinitely many descriptions of $\mathbf{S}^3 \times \mathbf{S}^2$ and $\mathbf{S}^3 \tilde{\times} \mathbf{S}^2$ as biquotients which do not admit a free (isometric) \mathbf{S}^1 action.*

Proof. Let S^1 act on $S^3 \times S^3$ via (5.3) with $a = -1, b = 3, c = 15k + 1$ and $d = 5, k \in \mathbf{Z}$. The action is free, since $\gcd(ab, cd) = 1$. Since the second Stiefel-Whitney class of the quotient $M_{a,b,c,d}^5$ is $w_2(M_{a,b,c,d}^5) = a + b + c + d = k \in \mathbf{Z}_2$, it therefore follows that both $S^3 \times S^2$ and $S^3 \tilde{\times} S^2$ arise, depending on the parity of k . Finally, we note that equation (8.1) can never be satisfied for any value of k . \square

REFERENCES

- [Ba] D. Barden, *Simply connected five-manifolds*, Ann. Math. **82** (3) (1965), 365–385
- [BB] L. Bérard-Bergery, *Les variétés riemanniennes homogènes simplement connexes de dimension impaire à courbure strictement positive*, J. Math. pure et appl. **55** (1976), 47–68
- [Br] G. E. Bredon, *Introduction to compact transformation groups*, Academic Press, New York, 1972
- [Ch] J. Cheeger, *Some examples of manifolds of non-negative curvature*, J. Diff. Geom. **8** (1973), 623–628
- [DV] J. DeVito, *The classification of simply connected biquotients of dimension at most 7 and 3 new examples of almost positively curved manifolds*, Ph.D. thesis (2011), University of Pennsylvania, <http://repository.upenn.edu/edissertations/311>
- [DL] H. Duan and C. Liang, *Circle bundles over 4-manifolds*, Arch. Math. **85** (2005), 278–282
- [Es] J.-H. Eschenburg, *Freie isometrische Aktionen auf kompakten Liegruppen mit positiv gekrümmten Orbiträumen*, Schriftenreihe Math. Inst. Univ. Münster (2) **32** (1984)
- [Fi1] R. Fintushel, *Circle actions on simply connected 4-manifolds*, Trans. Amer. Math. Soc. **230** (1977), 147–171
- [Fi2] R. Fintushel, *Classification of circle actions on 4-manifolds*, Trans. Amer. Math. Soc. **242** (1978), 377–390
- [GG] F. Galaz-Garcia, *Nonnegatively curved fixed point homogeneous manifolds in low dimensions*, Geom. Dedicata, to appear.
- [GGS] F. Galaz-Garcia and C. Searle, *Low-dimensional manifolds with non-negative curvature and maximal symmetry rank*, Proc. Amer. Math. Soc. **139** (2011), 2559–2564
- [Gr] K. Grove, *Geometry of, and via, symmetries*, Conformal, Riemannian and Lagrangian geometry (Knoxville, TN, 2000), Univ. Lecture Ser., vol. 27, Amer. Math. Soc., Providence, RI, 2002, 31–53
- [GS1] K. Grove and C. Searle, *Positively curved manifolds with maximal symmetry-rank*, J. Pure Appl. Algebra **91** (1994), no. 1-3, 137–142
- [GS2] K. Grove and C. Searle, *Differential topological restrictions curvature and symmetry*, J. Diff. Geom. **47** (1997), 530–559
- [GW] K. Grove and B. Wilking, *A knot characterization and 1-connected nonnegatively curved 4-manifolds with circle symmetry*, preprint 2011
- [GWZ] K. Grove, B. Wilking and W. Ziller, *Positively curved cohomogeneity one manifolds and 3-Sasakian geometry*, J. Diff. Geom. **78** (2008), no. 1, 33–111
- [GZ] K. Grove and W. Ziller, *Lifting group actions and nonnegative curvature*, Trans. Amer. Math. Soc. **363** (2011), no. 6, 2865–2890
- [HK] W. Y. Hsiang and B. Kleiner, *On the topology of positively curved 4-manifolds with symmetry*, J. Differential Geom. **29** (1989), no. 3, 615–621
- [Ke] M. Kerin, *On the curvature of biquotients*, Math. Ann., to appear
- [Kl] B. Kleiner, *Riemannian four-manifolds with non-negative curvature and continuous symmetry*, Ph.D. thesis, University of California, Berkeley, 1990
- [KMP] S. Kim, D. McGavran and J. Pak, *Torus group actions in simply connected manifolds*, Pacific J. Math. **53** (1974), no. 2, 435–444
- [Oh1] H. S. Oh, *6-dimensional manifolds with effective T^4 actions*, Topology Appl. **13** (1982), no. 2, 137–154
- [Oh2] H. S. Oh, *Toral actions on 5-manifolds*, Trans. Amer. Math. Soc. **278** (1983), no. 1, 233–252
- [Or] P. Orlik, *Seifert manifolds*, Lecture Notes in Mathematics, Vol. 291. Springer-Verlag, Berlin-New York, 1972
- [OR1] P. Orlik and F. Raymond, *Actions of the torus on 4-manifolds. I*, Trans. Amer. Math. Soc. **152** (1970), 531–559
- [Pav] A. V. Pavlov, *Five-dimensional biquotients of Lie groups*, (Russian) Sibirsk. Mat. Zh. **45** (2004), no. 6, 1323–1328; translation in Siberian Math. J. **45** (2004), no. 6, 1080–1083
- [Pao] P. S. Pao, *Nonlinear circle actions on the 4-sphere and twisting spun knots*, Topology **17** (1978), no. 3, 291–296
- [Pe1] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, preprint 2002, arXiv:math/0211159v1 [math.DG]
- [Pe2] G. Perelman, *Ricci flow with surgery on three-manifolds*, preprint 2003, arXiv:math/0303109v1 [math.DG]

- [SY] C. Searle and D. Yang, *On the topology of non-negatively curved simply connected 4-manifolds with continuous symmetry*, Duke Math. J. **74** (1994), no. 2, 547–556
- [To] B. Totaro, *Cheeger manifolds and the classification of biquotients*, J. Diff. Geom. **61** (2002), 397–451
- [Ve] L. Verdiani, *Cohomogeneity one manifolds of even dimension with strictly positive sectional curvature*, J. Diff. Geom. **68** (2004), 31–72
- [Wa] N. Wallach, *Compact homogeneous Riemannian manifolds with strictly positive curvature*, Ann. of Math. **96** (1972), 277–295
- [Wi1] B. Wilking, *Torus actions on manifolds of positive sectional curvature*, Acta Math. **191** (2003), no. 2, 259–297
- [Wi2] B. Wilking, *Positively curved manifolds with symmetry*, Ann. of Math. **163** (2006), no. 2, 607–668
- [Wi3] B. Wilking, *Nonnegatively and positively curved manifolds*, Surv. Differ. Geom., Vol. 11, Int. Press, Somerville, MA, 2007

MATHEMATISCHES INSTITUT, EINSTEINSTR. 62, 48149 MÜNSTER, GERMANY

E-mail address: f.galaz-garcia@uni-muenster.de

E-mail address: m.kerin@math.uni-muenster.de